

## $^{40}\text{Ar}/^{39}\text{Ar}$ Dating of the Shaxi Porphyry Cu-Au Deposit in the Southern Tan-Lu Fault Zone, Anhui Province

YANG Xiaoyong<sup>1,2</sup>, ZHENG Yongfei<sup>1</sup>, XIAO Yilin<sup>1</sup>, DU Jianguo<sup>3</sup> and SUN Weidong<sup>2</sup>

1 CAS Key Laboratory of Crust-Mantle Materials and Environments, School of Earth and Space Sciences, University of Science and Technology of China, Hefei, Anhui 230026

2 CAS Key Laboratory of Isotope Geochronology and Geochemistry, Guangzhou Institute of Geochemistry, Guangzhou, Guangdong 510640

3 Anhui Institute of Geological Survey, Hefei, Anhui 230001

**Abstract:** Four samples of plagioclase and biotite from the Shaxi porphyry in the lower part of the Yangtze metallogenic belt were analyzed for age determination with the  $^{40}\text{Ar}/^{39}\text{Ar}$  method. The results yield reproducible ages of 126 Ma to 135 Ma with a high level of confidence according to the agreement between isochron and plateau ages. The four Ar-Ar ages are relatively consistent within the analytical error. These ages are also consistent with, but more precise than, previous K-Ar and Rb-Sr ages and thus provide better constraints on the time of porphyry formation and associated Cu-Au mineralization along the middle to lower part of the Yangtze metallogenic belt. The ages of 126 to 135 Ma are interpreted to represent the intrusive time of the Shaxi porphyry, so that the Cu-Au mineralization should have occurred later due to the post-magmatic hydrothermal event.

**Key words:** Shaxi porphyry-type Cu-Au deposit,  $^{40}\text{Ar}/^{39}\text{Ar}$  dating, Yangtze metallogenic belt, Tancheng-Lujiang fault belt, East China

### 1 Introduction

The middle-lower part of the Yangtze Valley is one of the most important areas for Jurassic-Cretaceous poly-Cu-Fe-Au deposits in China (Chang et al., 1991; Zhai et al., 1996; Pan and Dong, 1999). Although porphyry Cu-(Mo-Au) deposits are the world's primary source of these metals, porphyry Cu deposits in this poly-metallogenic belt are a major source of Cu. The Shaxi porphyry Cu-Au deposit is one of the important discoveries during the 1970s. It is located in the northwestern Luzong volcanic basin, the central part of Anhui Province, east-central China, where the Tan-Lu (Tancheng-Lujiang) fault passes through the ore district. According to Chang et al. (1991) and Zhai et al. (1996), the eastern part of the Yangtze Craton in eastern China is an important Fe-Cu metallogenic province, whose metallogeny was controlled by faults and aulacogens in the continental plate during the Jurassic. The dominant WNW and E-W trending deep fault system controlled the distributions of Cu mineralization. A series of igneous rocks in this region can be grouped into two associations according to their relationship to the metal genesis. The first one is a Fe-

related group, whereas the second one is a Cu-related group.

This paper is focused on the second group in order to understand the relationship between its petrogenesis and Cu mineralization, particularly age determination of porphyrite in association with the Shaxi Cu-Au deposit. Previous geochronological studies of igneous rocks in the Lower Yangtze Valley, including K-Ar and Rb-Sr dating, suggest that different rock associations were formed in the same time span about 160 to 105 Ma (Bureau of Geology and Mineral Resources of Anhui Province, 1987; Chang et al., 1991; Zhao et al., 2004; Di et al., 2005; Xu et al., 2005). This study provides more accurate Ar-Ar ages from fast neutron activation techniques for biotite and plagioclase separates. The results are important for understanding the forming processes of both intrusives and Cu-Au deposits in the region.

### 2 Geological Setting

The Shaxi porphyrite intrusive is a small intrusion with an outcrop area of approximately 1 km<sup>2</sup>, but it is heavily copper mineralized and accompanied by Au mineralization (Yang, 1996; Yang et al., 2001b, 2002,

\* Corresponding author. E-mail: xyyang@ustc.edu.cn.




Fig. 1. The regional geological and tectonic map of the studied area.

(a) The regional geological and tectonic map along the middle-lower Yangtze metallogenic belt (after Chang et al., 1991).

SX – Shaxi-Changpushan porphyry Cu-Au deposit; AQ – Anqing magmatic Cu-Mo ore field; TL – Tongling skarn Cu-Au ore field.

(b) geological map of the Shaxi porphyry copper (gold) deposit, Central Anhui (after the No. 327 Geological Party, 1982).

1. Quaternary system; 2. red sedimentary rocks of the Kz-Cz series; 3. volcanic rocks of Kz; 4. clastic rocks of Pz; 5. clastic rocks of Pz; 6. middle and deep facies of diorite; 7. hypabyssal porphyry dioritoid; 8. ore-bearing quartz porphyry diorite; 9. subvolcanic rocks; 10. intrusive breccias; 11. explosion breccias; 12. extrapolated axis of anticline; 13. secondary axis of syncline; 14. fault; 15. region with drilling holes (including ZK3301 and ZK3307); 16. range of deposits: I - the Shaxi porphyry copper (gold) ore deposit; II - the range of a prospecting area of porphyry copper-gold deposit with similar mineralization of that in Shaxi (after Yang et al., 2001b).

2006; Yang and Lee, 2005). Figure 1a shows the general information about the regional tectonic setting along the middle-lower Yangtze metallogenic belt, and Fig. 1a is a geological map of the Shaxi porphyry copper (gold) deposit.

The studied region is located on the northern margin of the Yangtze Craton (Ren, 1980), but still south of the Dabie ultrahigh-pressure (UHP) metamorphic belt (Zhao et al., 2004a). The Dabie orogen was formed by Triassic subduction of the South China Block beneath the North China Block to result in continental collision (e.g., Zheng et al., 2003, 2005). The Dabie UHP metamorphic complex is truncated at its eastern end by the Tanlu fault, which offsets it by approximately 500 km northward to the Shandong Peninsular where it occurs as the Sulu orogen. Many studies have contributed to understanding of the geodynamics of the continental subduction and subsequent exhumation from various aspects of field geology, structural geology, mineralogy, petrology, geochemistry, geophysics, petrophysics and tectonics in the Dabie-Sulu orogenic belt (Zheng et al., 2003, 2005 and references therein). Zircon U-Pb dating by the TIMS (thermal ion mass spectrometer, upper intercept age) and SIMS (ion

microprobe,  $^{206}\text{Pb}/^{238}\text{U}$  age for zircon cores) has yielded a range of 620 to 880 Ma (mode range of 700 to 800 Ma) for protoliths of most UHP eclogites and orthogneisses in the Dabie-Sulu orogenic belt, corresponding to bimodal magmatism along the northern margin of the South China Block in response to the middle Neoproterozoic rifting tectonics (Zheng et al., 2004, 2006). Post-collisional igneous rocks are composed of voluminous coeval granitoids and minor mafic-ultramafic rocks that have emplacement ages of 117 to 135 Ma as established by different techniques of geochronology (e.g., Chen et al., 1995; Zhao et al., 2004b, 2005; Xie et al., 2006). Available results from element and isotope analyses indicate that the Early Cretaceous granitoids in the Dabie orogen have the geochemical characteristics similar to the subducted continent in many aspects. Therefore, they are petrogenetically considered as being generated by partial melting of the South China Block due to extension collapse of the collision-thickened orogen (Zhao et al., 2004b, 2005).

The whole district is affected by two multiple faults: the Tan-Lu fault which passes through the whole district, and the Fanshan-Tongling fault, which is also a regional deep

fault belt controlling the distribution and mechanism of the porphyry intrusive (Dong, 1984; Chang et al., 1991). The Tan-Lu fault belt, as a major wrench fault to the northwest of the Pacific Ocean, cuts the eastern continental crust of northeastern Asia and crosses more than several thousands kilometers in eastern China (Xu et al., 1987). Its northern part enters the Kurtuhin fault belt in the far east of Russia (Salon, 1977). Xu et al. (1987) presented the fact that this fault system would be traced for more than 3600 km, extending intermittently to South China, with a strike from NE to NNE, assuming an "S-shaped or snake-shaped" curve. This fault is 10–50 km in width and shows a fractural down-warping zone with highly plastic deformed ductile shear zones at levels of the middle to lower crust, and there are a series of ductile shear zones also with NE or NNE trends which were developed in the slide formation of the major fault belt within or near the studied region (Wang et al., 1983; Yang, 1992; Zhang, 1992; Yang et al., 1998a, b; 2001a).

### 3 Petrography and Geochemistry

#### 3.1 Petrography

Igneous rocks are widely distributed in the studied region with the characteristics of multiple magmatic activities. The period of the magmatic activity is clear from the field observation, where volcanic rocks in the deposit are overlying sandstones of the Xiangshan Group (middle-upper Jurassic) and have an intrusive contact with the lower Cretaceous red beds. Therefore, the volcanic activity was limited to the late Jurassic to early Cretaceous. The intrusive rocks in the Shaxi porphyry Cu-Au deposits are stocks, tongues, apophyses, ethmoliths and dikes, and include deep-shallow-ultra-shallow intrusive bodies. According to drilling and geophysical exploration (The No. 327 Geological Party, 1982; Yang et al., 2001b), all of the intrusive bodies have the same basement, and the ore bodies are resident in the upper part of the intrusive at 0–800 m depth. There are two active stages of igneous activity in this region. The early stage is an intrusive one, with different extents of Cu-Au mineralization, hosted mainly by quartz diorite porphyry and biotite quartz diorite porphyry with lesser amounts of amphibole diorite porphyry and breccia diorite porphyry. The late stage of magmatic activity is an injection accompanied with a large amount of basic, intermediate and intermediate-acid magmas with volcanic explosion and sub-volcano activity without any Cu-Au mineralization.

There are several kinds of intrusive rocks with porphyry copper mineralization in the Shaxi porphyry Cu-Au deposit. These are quartz diorite porphyry, biotite-quartz diorite porphyry and fine- to medium-grained porphyry diorite, which have a subhedral seriate texture.

Figure 2 shows the petrologic characteristics from observations of the intrusives from the Shaxi Cu-Au-bearing igneous intrusive in the lower part of the Yangtze metallogenic belt.

They contain phenocrysts of plagioclase and alkali-feldspar with sizes of 8–3 mm and 5–1.5 mm, respectively. Some feldspars are severely altered to sericite, chlorite and kaolinite in the alteration zones (Fig. 2a, b). The diorites in this region contain amphibole, quartz, microcline, biotite, muscovite, apatite, sphene, pyrite, magnetite and rare rutile. The microcline occurs as subhedral crystals with cross-hatch twinning of microperthitic and contains plagioclase. The biotite is all subhedral (Fig. 2c); however, the amount of biotite is more than that of muscovite in the diorite porphyry samples. All the opaque minerals are distributed among the main rock-forming minerals. The plagioclase in the diorite porphyry usually occurs as subhedral polysynthetically twinned crystals (Fig. 2d).

#### 3.2 Geochemistry

Late Mesozoic igneous rocks associated with Cu-Au mineralization crop out widely in the Lower Yangtze region. They intruded into either the Neoproterozoic low-grade metamorphic rocks or the Paleozoic to Triassic sedimentary strata.

##### 3.2.1 Major elements

Several rock associations were recognized in this region. The Luzong meso-volcanic rocks and some in the Tongling region with skarn Cu-Au mineralization fall into the shoshonite field in the  $K_2O$  vs.  $SiO_2$  plot (Fig. 3). The K-enriched rock association contains shoshonite series and ultrapotassic rocks classified by high  $Na_2O+K_2O$  of 8.1% to 12.0%, high  $K_2O$  content of 4.1% to 8.5%, and with  $K_2O/Na_2O$  ratios of about 0.8 to 1.4 (Bureau of Geology and Mineral Resources of Anhui Province, 1987; Wang and Yang, 1996; Yang, 1996; Xing and Xu, 1999; Yang and Lee, 2005). They are basaltic trachyandesite, trachyandesite and syenite in composition. However, the rocks in the Shaxi porphyry Cu-Au deposit are of calc-alkaline series, and the rocks in the Anqing and Tongling areas are of high-potassic calc-alkaline series. According to Chen et al. (2001), there exists a kind of rock series with Na-rich alkaline mafic association in this region showing low  $SiO_2$  of 46–56% and high alkali content with  $Na_2O/K_2O$  ranging from 5.0–7.1% and  $Na_2O/K_2O$  ratios of 1.4 to 4.3 (Xing and Xu, 1999). The associations of high potassic calc-alkaline series or calc-alkaline series occurring in the area north of the Yangtze River consisting of monzonite and granite stocks, diorite, quartz diorite and granodiorite stocks are distributed along the Yangtze River. These rocks are closely associated with the mineralization of copper, iron, sulfur and gold ore deposits




Fig. 2. Microscopic images of petrologic observation of the granitoids related to the Shaxi porphyry Cu-Au deposit.

(All the images are taken with the Laca microscope under the polarized light condition, the scale bar for each of the image is 0.20 mm. Abbreviations in the images: Am – amphibole; Bi – biotite; Mi – microcline; Pl – plagioclase; Chl – chlorite and chloritization; Py – pyrite; Q – quartz; Ser – sericite and sericitization).

1. Heavy alteration of sericitization and argillation in porphyrite from the Shaxi intrusive related to the porphyry Cu-Au deposit (ZK1405-21); 2. heavy alteration of sericitization and chloritization in porphyrite from the Shaxi intrusive (ZK1002-10); 3. regular shape of fresh biotite and plagioclase with zoning texture of porphyrite (Sample ZK 3303-07); 4. regular shape of fresh plagioclase with some smaller crystals of quartz and biotite, and one crystal of microcline can be identified from porphyrite ( Sample ZK 3303-01).

(Chang et al., 1991), making the area an important metal production base in China. Intrusions distributed in the region south of the Yangtze River include granites and granodiorites, occurring as batholiths and stocks. They are S-type granites in terms of chemical composition and mineralogy (Chang et al., 1991).

### 3.2.2 REE

The total (REE+Y) contents range from 256 ppm to 79 ppm (with an average value of 115.93 ppm) in the intrusive rocks in the Shaxi area (Yang, 1996). Almost all of the samples have the characteristics of enrichment in LREE and depletion in HREE without Eu anomaly compared to those of REE data from the other regions adjacent to Shaxi, such as the Anqing, Tongling, Luzong

and Chuxian areas (see Fig. 4). The REE patterns in the intrusive rocks from the Shaxi area are more similar to those of the lower crust rocks (Taylor et al., 1985).

### 3.2.3 Trace elements

The spidergram of trace elements in the Shaxi area is shown in Fig. 5, which has the characteristics of mantle compatible elements such as Sc, Cr, Co and Ni and some transitional compatible elements such as Ti, V, Mn, Fe and Cu, with strong fractionation compared to the average contents of trace elements of crust rocks and adjacent dioritoids along the Yangtze metallogenic belt (Thorpe, 1976; Taylor et al., 1985; Bureau of Geology and Mineral Resources of Anhui Province, 1987; Chang et al., 1991; Ren et al., 1991), especially those of Cu with very largely




Fig. 3.  $\text{SiO}_2$  vs.  $\text{Al}_2\text{O}_3$ ,  $\text{K}_2\text{O}$  and  $\text{Na}_2\text{O}/\text{K}_2\text{O}$  diagrams showing variations of different igneous rock associations related to Cu-Au mineralization in Anhui Province (after Rollinson, 1993 and Rickwood, 1989; data are derived from the No. 327 Geological Party, 1982; the No. 321 Geological Party, 1990; Bureau of Geology and Mineral Resources of Anhui Province, 1987; Chang et al., 1991; Ren et al., 1991; Qiu, 1992; Yang, 1996; Wang and Yang, 1996; Yang et al., 2001b).

positive anomaly, which may be interpreted as an important cause for Cu mineralization in this region. The lithophile elements such as K, Rb, Th, Sr, Ba, Li and La are also enriched compared to the average contents of the crustal rocks and adjacent dioritoids (Bureau of Geology and Mineral Resources of Anhui Province, 1987; Chang et al., 1991; Ren et al., 1991), which manifests the regional geochemical anomalies of these elements and may be interpreted as an important cause for Cu mineralization in this region. The hydrothermal alteration to some of the




Fig. 4. Normalized REE patterns of the altered intrusive rocks in the Shaxi porphyry Cu-Au deposit (data from Yang, 1996).

Cu-related intrusive rocks is another cause for the increase of these elements.

## 4 $^{40}\text{Ar}/^{39}\text{Ar}$ Dating

### 4.1 Sample selection and experimental methods

Biotite and plagioclase are commonly used for the  $^{40}\text{Ar}/^{39}\text{Ar}$  geochronometer. The fresh rocks selected for analysis in this study are porphyrites from the Shaxi intrusive. The rocks were crushed, separated using heavy liquid, and single biotite and plagioclase crystals were picked by hand under the binocular microscope to get pure monominerals mainly avoiding the intergrowths of minerals, fluid inclusions and alteration. Figure 2 (c, d) are the microscopic images of petrologic observation of the two selected porphyrite samples in the Shaxi porphyry Cu-Au deposit, showing the fresh minerals of both biotite and plagioclase suitable for the  $^{40}\text{Ar}/^{39}\text{Ar}$  geochronometer in this study.

It is well known that obtaining meaningful ages from such spectra is very difficult because of the possibility of alteration of minerals. Although biotite is a common rock-forming mineral suitable for Ar-Ar dating, it is often affected by alteration related to unique geological localities. The capability to obtain reliable ages depends on delicate choice and preparation of the fresh minerals, which will be the extremely important for obtaining the true geological dating. In this work great efforts were made, e.g., high-resolution binocular microscope was hired to check the pure biotite after heavy liquid separation in order to avoid the affects of alteration. In this way the purification of minerals was ensured for the further measurement, which is essential to the  $^{40}\text{Ar}/^{39}\text{Ar}$  geochronometer.

The preparation and conditions of the biotite and plagioclase samples and standard samples for analysis




Fig. 5. Spidergrams in the Shaxi porphyry copper (gold) deposit (data from Yang, 1996).  
(a) Transitional elements; (b) large lithophile elements.

with the fast neutron irradiation incremental heating for Ar extraction is described as follows: the separated monomineral samples were wrapped up in aluminum foil and placed in the central part of the B8 hole site of the 49-2 type reactor for fast irradiation with an irradiation time of 52 h 47 min, intermittent flux of neutrons  $6.63 \times 10^{12}$  n/cm<sup>2</sup>·s and integrated flux of neutrons  $1.26 \times 10^{18}$  n/cm<sup>2</sup>.

The standard samples used to monitor neutron fluxes included Chinese standard ones of ZBJ (hornblende) and ZBH-25 (biotite) with ages of  $132.8 \pm 1.4$  Ma and  $132.7 \pm 1.2$  Ma, respectively, and French B600 (biotite), Australia 77600 (hornblende) and international standard BSP (hornblende) with ages of 322 Ma,  $414.5 \pm 3.7$  Ma and  $2060 \pm 8$  Ma, respectively. The values of irradiation parameters  $J$  are 0.012826 (for sample ZK3303-07, biotite and plagioclase) and 0.012831 (for sample ZK3303-01, biotite and plagioclase), respectively.

The measurements were performed on an English RGA-10 Model gas-source mass spectrometer (VSS Company); the sample weights are from 0.14930 g to 0.1022 g. The decay constant  $\lambda$  is  $5.54 \times 10^{-10}/\text{a}$  in the static mode. The blanks of the whole system are  $^{40}\text{Ar} = 1.6 \times 10^{-14}$  mol and  $^{36}\text{Ar} = 1.2 \times 10^{-16}$  mol and the errors involved in the Ar measurement are within the range of 0.5%–1%.

#### 4.2 Results of the $^{40}\text{Ar}/^{39}\text{Ar}$ Ar dating

The gas released from the samples when heated in a tight container was analyzed on the mass spectrometer for Ar isotopes. Both biotite and plagioclase separates from two samples were analyzed and the results are shown in Table 1. Figures 6–9 show the apparent  $^{40}\text{Ar}/^{39}\text{Ar}$  age spectra and the isochron ages for the 4 selected samples.

The Ar isotope data acquired at 11 heating stages range from 420°C to 1450°C for each sample, though the

heating temperature steps were slightly different. The data points involved in plateau age calculation can satisfy such preconditions that the sample was formed under a chemically closed condition. The two biotite samples and two plagioclase samples represent ideal grains for  $^{40}\text{Ar}/^{39}\text{Ar}$  Ar dating, giving relatively reproducible ages with a high level of confidence ranging from 126 to 135 Ma according to the result of isochron age and plateau age analysis.

## 5 Discussion

It is desirable to identify apparent ages that are adversely influenced by alteration and to obtain meaningful ages from all rocks regardless of the conditions because many rocks are geologically unique and/or representative of key localities. However, it is difficult to obtain geologically meaningful ages from highly altered biotite using incremental step-heating. Even those biotites which appear pristine in thin sections can yield variable plateau and unexpected apparent ages (e.g., Lo and Onstott, 1989; Ruffet et al., 1991; Onstott et al., 1995; Roberts et al., 2001; Burgess et al., 2004). Biotite and plagioclase are the minerals most commonly dated by  $^{40}\text{Ar}/^{39}\text{Ar}$  geochronology. However, secondary alteration of biotite to phases such as chlorite is a common and widespread occurrence in nature and has long been known to lead to a lowering of biotite ages (Mitchell and Taka, 1984; Kelley et al., 1997; Adams and Kelley, 1998; Haines et al., 2004). Although later thermal disturbance can also yield plateau ages, alteration and weathering of grains are often correlated with  $^{40}\text{Ar}/^{39}\text{Ar}$  Ar age spectra that are seriously disturbed.

From the results of biotite and plagioclase, it can be

**Table 1 Data for  $^{40}\text{Ar}/^{39}\text{Ar}$  dating from the Shaxi porphyry intrusive, Anhui Province**

Heating stage	Temperature	$(^{40}\text{Ar}/^{39}\text{Ar})_m$	$(^{36}\text{Ar}/^{39}\text{Ar})_m$	$(^{37}\text{Ar}/^{39}\text{Ar})_m$	$(^{38}\text{Ar}/^{39}\text{Ar})_m$	$^{39}\text{Ar}_K \times 10^{-12}$ mol	$(^{40}\text{Ar}^*/^{39}\text{Ar}_K)$	$^{39}\text{Ar}_K$ $\pm 1\sigma$	$t$ (Ma) $\pm 1\sigma$
Sample ZK3303-07, biotite, $J = 0.012826$ , weight = 108.1 mg									
1	400	24.244	0.0431	0.1112	0.0503	3.244	11.55±0.472	1.25	249.34±11.34
2	500	17.009	0.0287	0.0986	0.0430	4.848	8.572±0.023	1.88	188.19±4.70
3	600	10.082	0.0192	0.0638	0.0303	8.421	4.413±0.008	3.28	99.34±1.42
4	700	9.0322	0.0107	0.0692	0.0387	21.57	5.847±0.007	8.40	131.05±1.77
5	800	7.1621	0.0040	0.0330	0.0317	34.33	5.971±0.004	13.3	133.14±1.65
6	880	7.1631	0.0042	0.0303	0.0304	32.71	5.912±0.004	12.7	131.88±1.64
7	960	6.7105	0.0026	0.0247	0.0337	52.89	5.937±0.004	20.6	132.41±1.63
8	1040	6.6853	0.0028	0.0260	0.0365	41.29	5.860±0.004	16.0	130.76±1.62
9	1140	7.1818	0.0045	0.0351	0.0345	25.51	5.846±0.004	9.94	130.46±1.63
10	1300	7.6829	0.0060	0.0389	0.0402	19.02	5.892±0.005	7.41	131.44±1.68
11	1400	8.6727	0.0090	0.0421	0.0490	12.75	6.002±0.007	4.97	133.81±1.81
Sample ZK3303-07, plagioclase, $J = 0.012826$ , weight = 149.3 mg									
1	420	33.137	0.0686	0.68121	0.25098	2.365	13.02±0.091	1.11	278.76±23.79
2	540	21.208	0.0496	0.93266	0.13087	3.454	7.473±0.038	1.62	188.13±6.30
3	660	13.289	0.0326	0.64155	0.08950	7.911	3.763±0.016	3.34	85.05±1.68
4	780	9.9710	0.0145	0.32557	0.04246	16.00	5.734±0.008	7.53	128.03±1.83
5	880	7.9326	0.0076	0.29386	0.02830	24.12	5.692±0.005	11.3	127.13±1.61
6	960	7.2052	0.0052	0.33906	0.02218	35.02	5.672±0.004	16.4	126.70±1.56
7	1040	7.1739	0.0050	0.42779	0.02210	32.00	5.714±0.004	15.0	127.61±1.57
8	1120	6.9550	0.0044	0.45492	0.02219	41.28	5.668±0.004	19.4	126.61±1.55
9	1200	7.0317	0.0047	0.71731	0.02740	29.21	5.668±0.004	13.7	127.04±1.57
10	1320	8.9710	0.0115	0.93129	0.03980	15.99	5.636±0.007	7.52	125.92±1.70
11	1450	13.320	0.0234	0.75508	0.08398	5.936	6.491±0.016	2.79	144.28±2.79
Sample ZK3301-01, biotite, $J = 0.012831$ , weight = 102.2 mg									
1	420	17.130	0.0257	0.1409	0.0699	5.405	9.542±0.024	2.32	208.37±5.32
2	540	10.240	0.0219	0.0803	0.0599	10.55	3.784±0.009	4.53	85.54±1.30
3	660	8.5542	0.0084	0.0421	0.0413	19.25	6.076±0.006	8.26	135.45±1.79
4	780	7.7454	0.0054	0.0264	0.0267	25.51	6.141±0.005	10.9	136.84±1.72
5	880	8.1081	0.0067	0.0397	0.0322	34.33	6.122±0.005	14.7	136.44±1.75
6	960	8.6725	0.0088	0.0548	0.0433	26.21	6.073±0.006	11.2	135.39±1.81
7	1040	7.5254	0.0050	0.0399	0.0307	27.37	6.031±0.005	11.7	134.49±1.69
8	1140	7.7325	0.0058	0.0440	0.0315	39.90	6.024±0.005	17.1	134.34±1.70
9	1250	8.7790	0.0093	0.0480	0.0451	19.95	6.046±0.007	8.56	134.81±1.81
10	1350	9.5294	0.0117	0.0553	0.0460	15.77	6.073±0.008	6.77	135.38±1.89
11	1450	12.486	0.0216	0.0735	0.0735	8.583	6.136±0.014	3.68	136.73±2.44
Sample ZK3301-01, plagioclase, $J = 0.012831$ , weight = 138.3 mg									
1	420	26.878	0.0578	0.60524	0.0953	4.011	9.929±0.058	2.38	216.35±12.17
2	540	10.107	0.0214	0.47937	0.0495	8.650	3.863±0.009	5.14	86.70±1.28
3	660	9.3469	0.0122	0.46373	0.0365	11.36	5.787±0.007	6.76	129.24±1.77
4	780	8.4848	0.0090	0.32217	0.0277	15.30	5.847±0.006	9.10	130.51±1.69
5	880	8.0000	0.0076	0.42457	0.0204	24.35	5.782±0.005	14.4	129.13±1.63
6	960	8.4651	0.0093	0.47780	0.0204	19.94	5.761±0.005	11.8	128.66±1.66
7	1020	7.7966	0.0068	0.54255	0.0199	27.36	5.839±0.004	16.2	130.34±1.63
8	1100	8.2291	0.0083	0.61051	0.0203	22.26	5.819±0.005	13.2	129.93±1.66
9	1200	8.6301	0.0095	0.79425	0.0224	16.92	5.857±0.006	10.0	130.73±1.70
10	1300	10.625	0.0166	0.93129	0.0302	11.12	5.785±0.009	6.62	129.19±1.89
11	1420	11.862	0.0206	0.77766	0.0458	6.742	5.839±0.011	4.0	130.35±2.12




Fig. 6. The  $^{40}\text{Ar}/^{39}\text{Ar}$  spectra and corresponding isochron diagrams of biotite in the Shaxi porphyrite intrusive, the southern Tan-Lu fault belt (Sample 3303-07).

(a) Spectral diagram; (b) isochron diagram.




Fig. 7. The  $^{40}\text{Ar}/^{39}\text{Ar}$  spectra (a) and corresponding isochron diagrams of plagioclase in Shaxi porphyrite intrusive, the south Tan-Lu fault belt (Sample 3303-07).

(a) Spectral diagram; (b) isochron diagram.

judged that these Ar-Ar ages can well represent the timing of the Shaxi porphyry emplacement, which can also be compared to the other ages by Ar-Ar, Rb/Sr and zircon U-Pb dating for many intrusives along the middle-lower Yangtze valley ranging from 140 to 122 Ma (Chen et al., 1985a, b, 1991, 1993, 1994; Zhou et al., 1987, 1988; McKee, 1988; Xu and Xing, 1994; Zhao et al., 2004a; Yang, 2006). The post-collisional magmatism in the Dabie orogen also occurred at 120 to 130 Ma (Chen et al., 1995; Zhao et al., 2004b, 2005; Xie et al., 2006). It is known that there is widespread occurrence of Early Cretaceous mineralizations in eastern China, but, significantly different settings of geotectonics are responsible for the same period of magmatism. Therefore, it remains to be resolved whether the Early Cretaceous magmatism along the middle-lower Yangtze Valley was triggered by mantle superwelling in response to the Pacific Superplume event (Zhao et al., 2004b, 2005) or it is related to subduction of the Pacific plate (Wu et al., 2005).

Recently, Xu et al. (2005) studied the Tongling skarn copper mineralized region and dated the main orebodies of the Dongguashan copper deposit with the Rb-Sr isotopic method, and they obtained the isochron ages of  $136.5 \pm 1.4$  Ma in the Qingshanjiao intrusive and  $134 \pm 11$  Ma of the later mineralized fluid inclusions in quartz vein. Di et al. (2005) studied the crystallization ages of the Xiaotongguanshan quartz monzodiorite and Shatanjiao quartz monzonite porphyry in the Tongling area using the SHRIMP zircon U-Pb method, and found that the crystallization age of the former is  $142.8 \pm 1.8$  Ma, that of the latter is  $151.8 \pm 2.6$  Ma. All the geochronology of the igneous rocks along the middle-lower Yangtze region indicates that these intrusives related to Cu-Au mineralization belong to the Late Jurassic-Early Cretaceous. It is believable that the ages from 126 to 135 Ma can represent the intrusive time in the Shaxi region, and the Cu-Au mineralization should have occurred later by a strong hydrothermal event of magmatism. These ages are also in accordance to the study of the movement of the Tan-Lu fault belt obtained by Ar-Ar dating such as Zhu et al. (2004), whose results of the  $^{40}\text{Ar}/^{39}\text{Ar}$  ages of mylonite whole-rock and muscovite from the later Tanlu ductile shear zone suggest another sinistral strike-slip cooling event at 128 Ma. Thus the strike-slip faulting of the Tan-Lu fault belt would induce a large-scale intrusion and doming uplift within its adjacent Shaxi area. Such a case was studied by Wu et al. (2004) in the Tongling region, and they provided a mechanism of emplacement in the deep and found the fact that the left-lateral shear shown by the ductile shear zone and the rheomorphic fold reveals that the pluton emplacement and the deformation of surrounding rock are controlled by a NNE-striking left-lateral shear stress field.






Fig. 8. The  $^{40}\text{Ar}$ - $^{39}\text{Ar}$  spectra and corresponding isochron diagrams of biotite in the Shaxi porphyrite intrusive, the south Tan-Lu fault belt (Sample 3301-01).

(a) Spectral diagram; (b) isochron diagram.




Fig. 9. The  $^{40}\text{Ar}$ - $^{39}\text{Ar}$  spectra and corresponding isochron diagrams of plagioclase in the Shaxi porphyrite intrusive, the south Tan-Lu fault belt (Sample 3301-01).

(a) Spectral diagram; (b) isochron diagram.

## 6 Conclusion

The Cu-Au mineralization in the Shaxi porphyry deposit occurs in a complex intrusive composed of quartz diorite porphyry, biotite-quartz diorite porphyry and fine- to medium-grained porphyry diorite along the lower part of the Yangtze metallogenic belt. The four reproducible Ar-Ar ages ranging from 126 to 135 Ma with a high level of confidence represent the formation ages of the Shaxi porphyry intrusive with Cu-Au mineralization. The ages constraining on the Cu-Au mineralization in the central Anhui Province are consistent with those of the majority of the adjacent acid intrusives with mass Cu-Au mineralization along the Yangtze metallogenic belt in the Yanshanian Period (Mesozoic).

## Acknowledgements

This study is supported by funds from the Ministry of Science and Technology of China (2006CB403500) and National Natural Science Foundation of China (40473021). Great thanks are due to Profs. Chang Yinbo, Tu Guangchi, Li Chaoyang, Wang Kuiren, Mao Jingwen and Yang Xueming for their guidance during our research. The No. 327 Geological Party of Anhui Province generously provided some samples and permission to access to the explored geological materials, and Sang Haiqing analyzed the Ar-Ar ages, which helped us greatly in the study.

Manuscript received Dec. 13, 2005

accepted Oct. 8, 2006

edited by Zhu Xiling

## References

- Adams, C.J., and Kelley, S., 1998. Provenance of Permian-Triassic and Ordovician metagraywacke terranes in New Zealand: Evidence from  $^{40}\text{Ar}/^{39}\text{Ar}$  dating of detrital micas. *Geological Society of American Bulletin*, 110: 422-432.
- Bureau of Geology and Mineral Resources of Anhui Province, 1987. *Regional Geology of Anhui Province* (Geological Memoirs, SW. 1, No. 5). Beijing: Geological Publishing House, 721 (in Chinese with English abstract).
- Burgess, R., Kiviets, G.B., and Harris J.W., 2004. Ar-Ar age determinations of eclogitic clinopyroxene and garnet inclusions in diamonds from the Venetia and Orapa kimberlites. *Lithos*, 77: 113-124.
- Chang Yinbo, Liu Xiangpei and Wu Yanchang, 1991. *The Copper-Iron Belt of the Middle and Lower Reaches of the Changjiang River*. Beijing: Geological Publishing House, 379 (in Chinese with English abstract).
- Chen, J.F., Foland, K.A., and Zhou, T.X., 1985a. Mesozoic granitoids of the Yangtze fold belt, China: Isotopic constrains on the magma sources. In: Wu, L.R., Yang, T.M., Yuan, K.R. et al. (eds.), *The Crust: The Significance of Granitic Gneisses in Lithosphere*. Athens: The Ophrastus, 217-237.
- Chen Jiangfeng, Li Xueming, Zhou Taixi and Foland, K.A., 1991.  $^{40}\text{Ar}/^{39}\text{Ar}$  dating of the Yueshan diorite, Anhui Province,

- and the estimated formation time of the associated ore deposit. *Geosciences*, 5: 91–99 (in Chinese with English abstract).
- Chen Jiangfeng, Xie Zhi, Liu Shunsheng, Li Xueming and Foland, K.A., 1995. Cooling age of Dabie orogen, China, determined by  $^{40}\text{Ar}/^{39}\text{Ar}$  and fission track techniques. *Sci. China (B)*, 38: 749–757.
- Chen, J.F., Yan, J., Xie, Z., Xu, X. and Xing, F.M., 2001. Nd and Sr isotopic compositions of igneous rocks from the Lower Yangtze Region in Eastern China: Constraints on Sources. *Physics and Chemistry of Earth (A)*, 26: 719–731.
- Chen Jiangfeng, Zhou Taixi and Foland, K.A., 1985b.  $^{40}\text{Ar}/^{39}\text{Ar}$  and Rb-Sr geochronology of the Qingyang batholith, Anhui Province, China. *Geochemica*, 4: 220–235 (in Chinese with English abstract).
- Chen Jiangfeng, Zhou Taixi, Li Xueming, Foland, K.A., Huang Chengyi and Lu Wei, 1993. Sr and Nd isotopic constraints on source regions of the intermediate and acidic intrusions from southern Anhui Province. *Geochemica*, 3: 261–268 (in Chinese with English abstract).
- Chen Jiangfeng, Zhou Taixi, Zhang Xun, Xing Fengming, Xu Xiang, Li Xueming and Xu Lihua, 1994. Isotopic geochemistry of Cu-mineralized intrusions from the Middle and Lower Reaches of Yangtze River. In: Chen, H.S. (ed.), *Isotope Geochemistry Studies*. Hangzhou: Zhejiang University Press, 214–229 (in Chinese).
- Dai Shengqian, Deng Jinfu, Wu Zongxu, Zhao Huailiang, Chen Jiangfeng and Du Jianguo, 2003. Evidence of magmatic petrology for Yanshanian orogeny in the Dabie orogen. *Geology in China*, 30: 159–165 (in Chinese with English abstract).
- Di Yongjun, Wu Ganguo, Zhang Da, Song Biao, Zang Wenshuan and Zhang Zhongyi, 2005. SHRIMP U-Pb zircon geochronology of the Xiaotongguanshan and Shatanjiao Intrusions and its petrological implications in the Tongling area, Anhui. *Acta Geologica Sinica* (English edition), 79(6): 795–802.
- Dong Shuwen, 1984. Identification of brush structure and its roll rock-and ore-controlling at Shaxi. *Acta Geologica Sinica* (Chinese edition), 58(3): 305–315 (in Chinese with English abstract).
- Haines, W., Turner, S., Kelley, S., Wartho, J.A and Sherlock, S.C., 2004.  $^{40}\text{Ar}/^{39}\text{Ar}$  dating of detrital muscovite in provenance investigations, a case study from the Adelaide Rift Complex, South Australia. *Earth Planet. Sci. Lett.*, 227: 297–311.
- Kelley, S., Bartlett, J.M., and Harris N.B.W., 1997. Pre-metamorphic Ar-Ar ages from biotite inclusions in garnet. *Geochim. Cosmochim. Acta*, 61: 3873–3878.
- Lo, C.H., and Onstott, T.C., 1989.  $^{39}\text{Ar}$  recoil artefacts in chloritized biotite. *Geochim. Cosmochim. Acta*, 53: 2697–2711.
- Mitchell, J.G., and Taka, A.S., 1984. Potassium and argon loss patterns in weathered micas: Implications for detrital mineral studies, with particular reference to the Triassic palaeogeography of the British Isles. *Sedimentary Geology*, 39: 27–52.
- McKee, E.H., 1988. K-Ar chronology of the Luohe iron district, Anhui Province, China. *Economic Geology*, 83: 433–435.
- No. 321 Geological Party, 1990. *A Comprehensive Prospecting on the Multiple Metallic Deposits in Tongling Area, South Anhui Province*. Hefei: Anhui Bureau of Geology and Mineral Resources, 136 (in Chinese).
- No. 327 Geological Party, 1982. *A Report of Geological Survey on Shaxi Porphyry Copper Deposit*. Hefei: Anhui Bureau of Geology and Mineral Resources, 163 (in Chinese).
- Onstott, T.C., Miller, M.L., Ewing, R.C., Arnold, G.W., and Walsh, D.S., 1995. Recoil refinements: Implications for the  $^{40}\text{Ar}/^{39}\text{Ar}$  dating technique. *Geochim. Cosmochim. Acta*, 59: 1821–1834.
- Pan Yuanming and Dong Ping, 1999. The Lower Changjiang (Yangzi/Yangtze River) metallogenic belt, east central China: Intrusion- and wallrock-hosted Cu-Fe-Au, Mo, Zn, Pb, Ag deposits. *Ore Geol. Rev.*, 15: 177–242.
- Qiu Ruilong, 1992. The petrological characteristics and genesis of the high potassium diorite in Yushan, Anqing, west Anhui Province. *Acta Petrologica Sinica*, 8: 97–108 (in Chinese with English abstract).
- Ren Jishun, Jiang Chuanfa, Zhang Zhengkun and Qin Deyu, 1980. *Geotectonic Evolution of China*. Beijing: Science Press, 1–124 (in Chinese).
- Ren Qijiang, Liu Xisheng and Xu Zhaowen, 1991. *Luzong Volcanic Basin of Mesozoic Era and the Mineralization, Anhui, East China*. Beijing: Geological Publishing House, 206 (in Chinese).
- Rickwood, C., 1989. Boundary lines within petrologic diagrams which use oxides for major and minor elements. *Lithos*, 22: 247–263.
- Rollinson, H.R., 1993. *Using Geochemical Data: Evaluation, Presentation, Interpretation*. New York: Longman Science and Technology, 48–51.
- Roberts, H.J., Kelley, S., and Dahl, S., 2001. Obtaining geologically meaningful  $^{40}\text{Ar}/^{39}\text{Ar}$  ages from altered biotite. *Chem. Geol.*, 172: 277–290.
- Ruffet, G., Feraud, G., and Amouric, M., 1991. Comparison of  $^{40}\text{Ar}/^{39}\text{Ar}$  conventional and laser dating of biotites from the North Tregor Batholith. *Geochim. Cosmochim. Acta*, 55: 1675–1688.
- Salon, S.A., 1977. Major tectonic characteristics and development history of the Sichote-Aline fold systems. *Geotectonics*, (1): 51–78 (in Russian).
- Sang Longkang, Wang Renjing, Zhang Zeming, Wang Qiang and He Bin, 2000. Late Yanshanian granites in the Jiuzhihe-Tiantangzhai area and uplift of core of the Dabie orogenic belt. *Acta Geologica Sinica* (Chinese edition), 74: 234–246 (in Chinese with English abstract).
- Taylor, S.R., and McLennan, S.M., 1985. *The Continental Crust: Its Composition and Evolution*. Oxford: Black Well Scientific Publications, 312.
- Thorpe, R.S., 1976. Rare earth data and petrogenesis of andesites from the north Chilean Andes. *Contrib. Mineral. Petrol.*, 54: 65–78.
- Wang Kuiren and Yang Xiaoyong, 1996. Geochemistry of Shaxi-Changpushan porphyry Cu-Au deposit from Anhui Province, East China. In: Pan Zhonghe et al. (eds.), *Advances in Solid Earth Sciences*. Beijing: Science Press, 154–164.
- Wang Xiaofeng, Wang Zhishun and Li Zhongjian, 1983. Study on the plastic deformation characteristics of the mylonites in the southern part of Tan-Lu fault belt. In: *Review in Structural Geology*. Beijing: Geological Publishing House, 30–36 (in Chinese with English abstract).
- Wu Ganguo, Zhang Da, Li Dongxu, Zhang Xiangxin, Shao Yongjun, Zang Wenshuan and Wang Qunfeng, 2004. Study of the emplacement mechanism of the Fenghuangshan granite pluton and related Cu-Au mineralization in Tongling, Anhui Province. *Acta Geologica Sinica* (English edition), 78(2): 492–502.
- Wu, F.Y., Lin, J.Q., Wilde, S.A., Zhang, X.O., and Yang, J.H., 2005. Nature and significance of the Early Cretaceous giant

- igneous event in eastern China. *Earth Planet. Sci. Lett.*, 233: 103–119.
- Xing Fengming and Xu Xiang, 1999. *Yangtze Magmatic Belt and Metallogenesis*. Hefei: Anhui People's Publishing House, 170 (in Chinese).
- Xie, Z., Zheng, Y.F., Zhao, Z.F., Wu, Y.B., Wang, Z.R., Chen, J.F., Liu, X.M., and Wu, F.Y., 2006. Mineral isotope evidence for the contemporaneous process of Mesozoic granite emplacement and gneiss metamorphism in the Dabie orogen. *Chem. Geol.*, 231: 214–235.
- Xu, J.W., Zhu, G., Tong, W.X., Cui, K.R., and Liu, Q., 1987. Formation and evolution of the Tancheng-Lujiang wrench fault system, a major shear system to the northwest of the Pacific Ocean. *Tectonophysics*, 134: 273–310.
- Xu Xiang and Xing Fengming, 1994. Whole-rock and mineral Rb-Sr isochron ages of the three gabbros in Nanjing-Wuhu area, China. *Scientia Geologica Sinica*, 29: 309–312 (in Chinese with English abstract).
- Xu Zhaowen, Lu Xiancai, Ling Hongfei, Lu Jianjun, Jiang Shoyong, Nie Guiping, Huang Shunsheng and Hua Ming, 2005. Metallogenetic Mechanism and timing of late superimposing fluid mineralization in the Dongguashan diplogenetic stratified copper deposit, Anhui Province. *Acta Geologica Sinica* (English edition), 79(3): 405–413.
- Yang Xiaoyong, 1992. *The Study of Tectonogeochemistry in Southern Part of Tan-Lu Fault Belt*. Thesis of Master Degree of the University of Science and Technology of China, 156 (in Chinese with English abstract).
- Yang Xiaoyong, 1996. *The Cu-Au Metallogenic Prospecting Areas from Middle-Lower Reaches of Changjiang River: A Study on Metallogenic Geochemistry of Some Typical Copper and Gold Ore Deposits*. Thesis of Ph.D. Degree of the University of Science and Technology of China, Hefei, 214 (in Chinese with English abstract).
- Yang Xiaoyong, 2006.  $^{40}\text{Ar}/^{39}\text{Ar}$  dating and geological significance on the Cu-bearing porphyrite of Shaxi from southern Tan-Lu fault belt. *J. Mineral. Petrol.*, 26(2): 52–56 (in Chinese with English abstract).
- Yang Xiaoyong, Liu Deliang, Yang Xueming, Wang Kuiren and Dai Jinxing, 1998a. Characteristics of compositional migration in mylonites from the ductile shear zones of the southern Tancheng-Lujiang fault belt, Eastern Anhui Province. *Acta Geologica Sinica* (English edition), 72(1): 37–50.
- Yang Xiaoyong, Yang Xueming, Liu Deliang, Wang Kuiren and Dai Jinxing, 1998b. A micro-structural evidences of tectonic superimposition (compound) by south part of Tancheng-Lujiang fault and Dabie-Jiangnan orogenic belt, China. *Acta Geophysica Sinica*, 41(sup): 123–132 (in Chinese with English abstract).
- Yang Xiaoyong, Liu Deliang and Wagner, G.A., 2001a. Conditions of deformation and variations of compositional and structural state of feldspars during mylonitization. Exemplified from the ductile shear zones in south Tancheng-Lujiang fault belt of China. *Neues Jahrbuch für Mineralogie (Mon.)*, J.g. (2001): 415–432.
- Yang Xiaoyong, Wang Kuiren, Sun Liguang, Yang Xueming, Li Yingxian and Shi Kunfa, 2001b. A prospecting porphyry Cu-Au deposit in Changpushan area by geochemical and geophysical exploration, central Anhui, east China. *Chime der Erder*, 61: 254–276.
- Yang Xiaoyong, Wang Kuiren, Yang Xueming and Sun Liguang, 2002. Characteristics of mineralization and gold occurrence in Shaxi porphyry copper-gold deposit, central Anhui, China. *Neues Jahrbuch für Mineral. (Abh)*, 177: 293–320.
- Yang Xiaoyong and Lee Insung, 2005. Geochemistry and Metallogenesis in Lower part of Yangtze metallogenic valley: A case study in Shaxi-Changpushan porphyry Cu-Au deposit and a review on the adjacent Cu-Au deposits. *Neues Jahrbuch Mineral. (Abh)*, 181: 223–243.
- Yang Xiaoyong, Yang Xueming, Jiang Laili, Wang Kuiren and Sun Liguang, 2006. Geochemical study of Shaxi porphyry copper-gold deposit in south part of Tan-Lu fault belt, East China. *J. Geol. Soc. India*, 67: 475–494.
- Zhai Yusheng, Xiong Yongliang, Yao Shuzhen and Lin Xinduo, 1996. Metallogeny of copper and iron deposits in the Eastern Yangtze Craton, east-central China. *Ore Geol. Rev.*, 11: 229–248.
- Zhang Jiasheng, 1992. The Characteristics and significance of Tan-Lu fault belt. *J. China Univ. Geosci.*, 18: 363–371 (in Chinese with English abstract).
- Zhao, Z.F., Zheng, Y.F., Wei, C.S., and Gong, B., 2004a. Temporal relationship between granite cooling and hydrothermal uranium mineralization at Dalongshan in China: a combined radiometric and oxygen isotopic study. *Ore Geol. Rev.*, 25: 221–236.
- Zhao, Z.F., Zheng, Y.F., Wei, C.S., and Wu, Y.B., 2004b. Zircon isotope evidence for recycling of subducted continental crust in post-collisional granitoids from the Dabie terrane in China. *Geophys. J. Res. Lett.*, 31, L22602, doi: 10.1029/2004GL021061.
- Zhao, Z.F., Zheng, Y.F., Wei, C.S., Wu, Y.B., Chen, F.K., and Jahn B.M., 2005. Zircon U-Pb age, element and C-O isotope geochemistry of post-collisional mafic-ultramafic rocks from the Dabie orogen in east-central China. *Lithos*, 83: 1–28.
- Zheng, Y.F., Fu, B., Gong, B., and Li L., 2003. Stable isotope geochemistry of ultrahigh pressure metamorphic rocks from the Dabie-Sulu orogen in China: Implications for geodynamics and fluid regime. *Earth-Sci. Rev.*, 62: 105–161.
- Zheng, Y.F., Wu Y.B., Chen F.K., Gong B., Li L., and Zhao Z.F., 2004. Zircon U-Pb and oxygen isotope evidence for a large-scale  $^{18}\text{O}$  depletion event in igneous rocks during the Neoproterozoic. *Geochim. Cosmochim. Acta*, 68: 4145–4165.
- Zheng, Y.F., Zhou, J.B., Wu, Y.B., and Xie, Z., 2005. Low-grade metamorphic rocks in the Dabie-Sulu orogenic belt: A passive-margin accretionary wedge deformed during continent subduction. *Int. Geol. Rev.*, 47: 851–871.
- Zheng, Y.F., Zhao, Z.F., Wu, Y.B., Zhang, S.B., Liu, X.M., and Wu, F.Y., 2006. Zircon U-Pb age, Hf and O isotope constraints on protolith origin of ultrahigh-pressure eclogite and gneiss in the Dabie orogen. *Chem. Geol.*, 231: 135–158.
- Zhou Taixi, Chen Jiangfeng, Li Xueming and Foland, K.A., 1988. Has the Indo-Sinian magmatism occurred in Anhui Province? *Acta Petrologica Sinica*, 4: 46–53 (in Chinese with English abstract).
- Zhou Taixi, Li Xueming, Zhao Junshen and Zhang Futao, 1987. Geochronology of igneous rocks from the Tongguanshan ore area of Anhui Province. *J. China Univ. Sci. Tech.*, 17: 403–407 (in Chinese with English abstract).
- Zhou Xunruo and Ren Jin, 1994. *The Mesozoic Granites along the Middle-Lower Part of Yangtze*. Beijing: Geological Publishing House, 1–119 (in Chinese with English abstract).
- Zhu Guang, Huo Mingjin, Wang Yongsheng, Liu Guosheng and Niu Manlan, 2004. Thermal evolution of the Tanlu fault zone on the eastern margin of the Dabie Mountains and its tectonic implications. *Acta Geologica Sinica* (English edition), 78(4): 940–950.