



Accelerated biodegradation of BPA in water-sediment microcosms with *Bacillus* sp. GZB and the associated bacterial community structure



Jukun Xiong^a, Taicheng An^b, Guiying Li^{b,*}, Ping'an Peng^a


^a State Key Laboratory of Organic Geochemistry, Guangzhou Institute of Geochemistry, Chinese Academy of Sciences, Guangzhou, 510640, China

^b Guangzhou Key Laboratory of Environmental Catalysis and Pollution Control, School of Environmental Science and Engineering, Institute of Environmental Health and Pollution Control, Guangdong University of Technology, Guangzhou, 510006, China

HIGHLIGHTS

- Bioaugmentation with GZB can effectively improve BPA biodegradation in sediments.
- Bioaugmentation was stimulated by adding electron donors/co-substrates.
- Bioaugmentation did not change the bacterial diversity.
- Bioaugmentation can change the composition of the major bacterial groups.

GRAPHICAL ABSTRACT



ARTICLE INFO

Article history:

Received 4 January 2017

Received in revised form

9 May 2017

Accepted 29 May 2017

Available online 31 May 2017

Handling Editor: Chang-Ping Yu

Keywords:

Bisphenol A
Bioaugmentation
Bacterial community
Microcosm

ABSTRACT

Bisphenol A (BPA) is a synthetic chemical primarily used to produce polycarbonate plastics and epoxy resins. Significant industrial and consumer's consumption of BPA-containing products has contributed to extensive contamination in different environmental matrices. In this study, microcosms bioaugmented with *Bacillus* sp. GZB were constructed to investigate BPA biodegradation, identify the main bacterial community, and evaluate bacterial community responses in the microcosms. Under aerobic conditions, BPA was quickly depleted as a result of bioaugmentation with *Bacillus* sp. GZB in water-sediment contaminated with pollutants. The pollutants used were generally associated with the electronic wastes (mobile phones, computers, televisions) dismantling process. Adding BPA affected the bacterial community composition in the water-sediment. Furthermore, BPA biodegradation was enhanced by adding electron donors/co-substrates: humic acid, NaCl, glucose, and yeast extract. Metagenomic analysis of the total 16S rRNA genes from the BPA-degrading microcosms with bioaugmentation illustrated that the genera *Bacillus*, *Thiobacillus*, *Phenylobacterium*, and *Cloacibacterium* were dominant after a 7-week incubation period. A consortium of microorganisms from different bacterial genera may be involved in BPA biodegradation in electronic waste contaminated water-sediment. This study provides new insights about BPA bioaugmentation and bacterial ecology in the BPA-degrading environment.

© 2017 Elsevier Ltd. All rights reserved.

1. Introduction

Bisphenol A (2,2-bis(4-hydroxyphenyl)propane, BPA, CAS No. 80-05-7) is widely used as an intermediate to produce

* Corresponding author.

E-mail address: ligy1999@gdut.edu.cn (G. Li).

polycarbonate, epoxy resins, flame retardants, and other products (Hoekstra and Simoneau, 2013). Worldwide demand and annual consumption of BPA continues to increase as a result of human activity (Huang et al., 2012). More than one million pounds of BPA are estimated to be released into the environment annually, mainly from the discharged plastic leachate in landfills, processing of BPA in manufacturing, and the combustion of computer printed circuit boards in electronic waste (Erler and Novak, 2010; USEPA, 2010; Huang et al., 2012; Peng et al., 2015). As a result, BPA has been detected in groundwater, industrial water, drinking water, municipal sewage sludge, and surface water (Belfroid et al., 2002; Huang et al., 2012; Careghini et al., 2015; Xiong et al., 2015; Toro-Velez et al., 2016). Further, BPA has been confirmed to be an endocrine-disrupting chemical, and may adversely affect endocrine functions in humans and wildlife. This may cause changes in developmental processes (Hutler Wolkowicz et al., 2016), altered reproductive capacity (Quan et al., 2017), damaged nervous systems (Beronius et al., 2013), increased cancers (Chen et al., 2002), and decreased immunocompetence (Lee et al., 2013). This has led to increased focus on the environmental fate of BPA (Sun et al., 2012; Toro-Velez et al., 2016).

The biological transformation of organics is one of the main transformation pathways in the environment, and microorganisms can effectively remove different environmental pollutants in the natural aquatic environment (Yang et al., 2016). However, most previous studies have focused on isolating different indigenous BPA-degrading bacterial species (Eio et al., 2014) and conducting BPA degradation tests using these bacteria in liquid culture (Wang et al., 2014). For example, BPA-degraders *Sphingomonas* sp. strain BP-7 (Yang et al., 2015), *Sphingomonas* sp. strain AO1 (Roh et al., 2009), *Achromobacter xylosoxidans* strain B-16 (Zhang et al., 2007), and *Bacillus* sp. GZB (Li et al., 2012) were isolated from different environments and their degradation efficiencies were studied in the laboratory culture. However, bacteria with strong biodegradation ability in the natural aquatic environments are limited and typically depend on biostimulation and/or bioaugmentation.

Bioaugmentation, using these isolated pollutant-degrading microorganisms, may be an attractive option to quickly remediate contaminated aquatic environments. For example, a tetrabromobisphenol A (TBBPA) degrader *Ochrobactrum* sp. T was used to bioaugment TBBPA degradation in river sediment; removal efficiencies were noticeably enhanced by bioaugmentation with this TBBPA-degrader (Li et al., 2016). However, little is known about bioaugmentation using BPA degraders in river sediment. In addition, it is unknown whether a bioaugmentation strategy could significantly change the whole bacterial community structure, such as when a BPA-degrader is added in the BPA-contaminated sediment. In addition, better understanding BPA-degrading bacterial communities can help researchers to better understand BPA bioaugmentation in natural aquatic environments, which can be useful for further practical application. This heightens the importance of studying the accelerated biodegradation of BPA in water-sediment microcosms when treated with *Bacillus* sp. GZB, and the resulting bacterial community structure.

Therefore, this study applied bioaugmentation with *Bacillus* sp. GZB, isolated from an electronic waste dismantling zone. Previous studies have demonstrated that bacteria could effectively degrade BPA in liquid culture. This study first investigated the biodegradation potential of BPA with bioaugmented with *Bacillus* sp. GZB in simulated water-sediment under aerobic conditions. Second, the study identified the dominant bacterial community responses to bioaugmentation using high-throughput sequencing. Third, we investigated the roles of different specific bacterial communities, based on enrichment during the bioaugmentation process. These

results can help optimize current remediation methods.

2. Materials and methods

2.1. Experimental design

Surface sediment samples were collected from a river (0–10 cm depth) in an electronic waste dismantling zone in Guangdong Province, South China. Samples were placed in sterilized glass bottles and stored at -20°C until they were analyzed. The sampling region was seriously polluted by heavy metals (Wu et al., 2015) and organics, including polycyclic aromatic hydrocarbon (PAHs) (Zhang et al., 2011), brominated flame retardants (BFRs), and BPA (Xiong et al., 2015). This contamination resulted from the primitive techniques used during electronic waste and plastic recycling. The inorganic salt medium used to isolate BPA-degrading bacteria and the growth medium were prepared based on previous research (Li et al., 2012, 2016). The pH value of the inorganic salt medium was adjusted to 7.0 before sterilization.

Microcosms were established using homogenized surface sediment, consisting of 47% (V/V) surface sediment, 53% (V/V) inorganic salt medium, and 10 mg L^{-1} of BPA (97%, Acros Organics). Each microcosm contained 150 mL homogenized surface sediment in a 250 mL flask. Six microcosms were established, each with a duplicate: (1) unamended controls under aerobic conditions; (2) aerobic conditions + bioaugmentation with *Bacillus* sp. GZB; (3) yeast extract (5 mg L^{-1}) + bioaugmentation with *Bacillus* sp. GZB; (4) NaCl (10 ng L^{-1}) + bioaugmentation with *Bacillus* sp. GZB; (5) humic acid (0.5 g L^{-1}) + bioaugmentation with *Bacillus* sp. GZB; and (6) glucose (10 mM) + bioaugmentation with *Bacillus* sp. GZB.

Bacillus sp. GZB was placed in sterilized growth medium and cultured at 37°C in a horizontal shaker at 200 rpm for 15 h. Then, 30 mL of the incubated growth medium was centrifuged and rinsed three times with stroke-physiological saline solution to collect the *Bacillus* sp. GZB. An identical wet weight (approximately 0.35 g) of *Bacillus* sp. GZB was added to each microcosm. These microcosms were incubated in a horizontal shaker (150 rpm) in the dark at 25°C . Previous research showed that the *Bacillus* sp. GZB isolated from an electronic waste recycling sediment from regions where electronic waste recycling is done could effectively degrade BPA under anaerobic and aerobic conditions (Li et al., 2012). Detailed sample collection methods can be found in previously published research (Li et al., 2016).

2.2. Chemical and molecular analyses

The residual BPA in water-sediment was extracted and measured based on existing references (Xiong et al., 2015, 2016). Genomic DNA of each water-sediment mixed samples was extracted using a E.Z.N.A.[®] Soil DNA Kit (Omega Bio-tek, Inc., USA). The bacterial V3-V4 region of the 16S rRNA gene was amplified using the forward primers 341F (5'-CCTACACGACGCTCTCCGATCTN-3') and reverse 805R (5'-GACTGGAGTTCCTTGGCACCCGAGAATTCCA-3') (Li et al., 2016; Xiong et al., 2017). The PCR programs for bacterial amplification ran on a Bio-rad T 100™ thermal cycler (California, USA) using previously published procedures (Li et al., 2016). PCR amplicons were evaluated using electrophoresis on 2.0% agarose gel and extracted with the SanPrep Column DNA Gel Extraction Kit (Sangon Biotech, China). The amplicons underwent high-throughput sequencing analysis (Illumina MiSeq sequencing) to characterize bacterial communities and examine their relative abundance and diversity.

The generated metagenomic sequences were filtered to check the quality control procedures; low quality sequences were removed based on reference (Li et al., 2016). UCLUST software




Fig. 1. Percentage of BPA residuals in water-surface sediment under different treatments.

(<http://www.drive5.com/uclust>) was used to assign sequences to operational taxonomic units (OTUs); sequences were clustered into OTUs at 97% sequence similarity. The identity of the representative sequence for each OTUs were selected based on the most abundant sequences; the taxonomy was assigned using the Ribosomal Database Project (RDP) classifier (Wang et al., 2007). Shannon diversity index and rarefaction curve of each sample were estimated from the sequence data using Mothur software. To compare the community diversity among samples based on genetic diversity, the Fast UniFrac online tool (<http://unifrac.colorado.edu/>) was used to estimate the weighted UniFrac metric; principal coordinate analysis (Pcoa) was then generated. Heat maps were constructed using the software tool Heml 1.0. The 16S rRNA gene sequences in this study were deposited in the National Center for Biotechnology Information (NCBI) Sequence Read Archive with accession numbers SRA308130.

2.3. Data analysis

Statistical analyses were conducted using Microsoft Excel 2010 and the Statistical Package for Social Sciences v18.0 software (SPSS Inc., IL, USA). Statistical significance between datasets was tested using analysis of one-way variance (ANOVA).

3. Results and discussion

3.1. BPA biodegradation

This study analyzed bioaugmentation and biostimulation processes in water-sediment microcosms. Fig. 1 illustrates the residual BPA patterns in the untreated control microcosms and in the microcosms with different treatments after a 7-week incubation period. During the incubation process, the degradation was negligible in the unamended control microcosms ($P > 0.05$). There was an average 35.9% BPA removal in the bioaugmented microcosms; the average BPA removal efficiencies were 56.3%, 44.4%, 47.6%, and

56.6% in bioaugmented microcosms with added yeast extract, NaCl, humic acid, and glucose, respectively. The differences between the unamended controls and the bioaugmented microcosms were statistically significant ($P < 0.01$).

These results confirmed that the bioaugmentation and biostimulation increased BPA removal in the water-sediment system. Furthermore, as expected, the BPA removal efficiency in the microcosms using *Bacillus* sp. GZB bioaugmentation was higher than the efficiencies in the untreated controls. This suggested that the bioaugmentation with *Bacillus* sp. GZB could enhance BPA biodegradation in water-sediment ($P < 0.01$). Therefore, the enhanced biodegradation efficiency was closely related to the added *Bacillus* sp. GZB, a facultative anaerobic BPA-degrading bacterium isolated from an electronic waste dismantling zone estuarine (Li et al., 2012).

Previous studies showed that xenobiotic biodegradation can be enhanced by adding electron donors/co-substrates (Li et al., 2016; Xiong et al., 2017). In this study, the data listed in Table 1 show that BPA degradation rates (k) increased in microcosms where bioaugmentation was supplemented with yeast extract, NaCl, humic acid, and glucose ($P > 0.05$). These results illustrated that these additional compounds could promote BPA bioaugmentation in the microcosms; this finding was consistent with existing literature (Xiong et al., 2017). Glucose donates H_2 and carbon during bacterial respiration, stimulating the bacteria's ability to degrade compounds (Zu et al., 2014). Yeast extract, a complex mixture of amino acids, peptides, and proteins (Fava et al., 1995), is often added as an additional alternative carbon source. It also protects the bacteria, reducing the adverse impact of the degradable compounds, and providing bacteria with good growth and biodegradation capacity. Humic acid effectively improves the sorption and binding of contaminated compounds onto sediment in contaminated aquifers (Conte et al., 2001). This facilitates contact between bacteria and BPA, facilitating BPA degradation. Salinity is another environmental factor that impacts the types of bacteria that colonize the sediment and their biodegradation potential (Xiong et al., 2017). In this study, BPA biodegradation was enhanced by adding 10 ng L^{-1} NaCl, suggesting that this salinity benefits BPA degradation.

3.2. Diversity of bacterial community

In this study, Illumina MiSeq sequencing analysis was used to investigate the water-sediment bacterial communities in the microcosm, using bioaugmentation at different incubation periods. Approximately 168,664 valid reads of 16S rRNA gene with an average length of approximately 455 bp were obtained after trimming and chimera removal. Using a 97% sequence similarity cutoff, the number of OTUs in the water-sediment bacterial communities varied from 1147 to 2861 (Table 2). These results indicated that the genera of bacterial community were highly diverse. The rarefaction curve was used to standardize and compare the observed taxon richness between samples, to identify whether the sample was unequally sampled (Xiong et al., 2017). In this study, the rarefaction curve for each water-sediment sample did not plateau, indicating a

Table 1
First order degradation rate constants (k) and half-lives ($t_{1/2}$) of BPA from the water-surface sediment in microcosms.

Treatments	$k \text{ (d}^{-1}) \pm \text{standard deviation}$	$t_{1/2} \text{ (d)}$	r^a
Bioaugmentation	$8.90 \times 10^{-3} \pm 0.85 \times 10^{-3}$	78	0.99
Yeast extract + bioaugmentation	$1.65 \times 10^{-2} \pm 0.49 \times 10^{-3}$	42	0.98
Sodium chloride + bioaugmentation	$1.19 \times 10^{-2} \pm 0.92 \times 10^{-3}$	58	0.99
Humic acid + bioaugmentation	$1.32 \times 10^{-2} \pm 0.28 \times 10^{-3}$	53	0.99
Glucose + bioaugmentation	$1.67 \times 10^{-2} \pm 0.35 \times 10^{-3}$	41	0.99

All degradation rate constants and half-lives are the mean of duplicate samples.

^a r : Correlation coefficient. Bioaugmentation: addition of *Bacillus* sp. GZB.

Table 2
Comparison of phylotype coverage and diversity estimators of the bacteria.

Sample	Reads ^a	OTUs ^b	Shannon	ACE	Chao1	Coverage
Week 0	10,131	1147	3.68	5086	3199	92.81
Week 1	10,417	1219	4.50	4821	3052	90.79
Week 2	12,254	2665	5.86	12,946	7699	86.52
Week 3	10,118	2735	5.93	13,660	8192	87.98
Week 4	13,993	2056	5.88	6269	4282	93.31
Week 5	23,611	1953	5.93	8778	5301	84.33
Week 7	88,140	2861	5.94	13,229	7831	87.88

^a Reads after filtering, trimming and normalizing.

^b The operational taxonomic units (OTUs) were defined with 97% similarity.




Fig. 2. Principal coordinate analysis (PcoA) of the samples using Weighted-UniFrac from pyrosequencing.

significantly higher water-sediment bacterial diversity at the sequencing depth used in this study (Fig. S1).

Shannon index values were used to assess bacterial community diversity (Xiong et al., 2017). In this study, the Shannon index values of the water-sediment samples increased from 3.68 to 5.94 during the incubation period. The results indicated a high diversity in bacterial 16S rRNA libraries during the BPA biodegradation process. The species richness indices of the abundance-based coverage estimator (ACE) and Chao1 varied from 5086 to 13,229, and from 3199 to 7831, respectively. These results also highlighted high bacterial species richness. In addition, calculated coverage, ranging from 86.52% to 93.31%, revealed that most bacterial communities were accounted for using 454 pyrosequencing. Principal coordinate analysis (PcoA) (Fig. 2) showed that water-sediment samples occupied the divergent positions. The diversity estimator indexes above suggested that bacterial community diversities did not decline in the microcosm bioaugmented with *Bacillus* sp. GZB during the incubation period. This suggested that bioaugmentation and xenobiotics (BPA) did not change bacterial diversity.

3.3. Composition of bacterial community

First, bacterial community composition was analyzed at the phylum level in the clone library. In this study, there were nine phyla with reads of more than 1.0% in the microcosm samples with bioaugmentation. The nine primary bacterial phyla in the sediment were *Proteobacteria*, *Firmicutes*, *Bacteroidetes*, *Actinobacteria*, *Acidobacteria*, *Verrucomicrobia*, *Planctomycetes*, *Chloroflexi*, and *TM7* (Fig. 3).

For the water-sediment in the microcosm, *Proteobacteria*




Fig. 3. Phylogenetic classification of the bacterial communities using pyrosequencing at the phylum level obtained from Ribosomal Database Project classifier analysis. The phyla with reads of less than 1.0% and unclassified phyla were grouped as "other" phyla.




Fig. 4. The taxonomic distribution of the *Proteobacteria* phyla for pyrosequencing.

(accounting for 37.5–59.1% of total bacteria) was the first largest phylum group in all samples; it mainly consisted of *Alpha-proteobacteria*, *Beta-proteobacteria*, and *Gamma-proteobacteria* (Fig. 4). Previous studies show that this *proteobacteria* group is widely distributed in waste water, sediment, and soil, where BPA levels are very high (Conte et al., 2001; Yang et al., 2015; Xiong et al., 2017). For example, Yang et al. found that BPA could be quickly depleted in BPA-spiked sediment; the microbial community structure in river sediment shifted, such that *Gamma-proteobacteria* and *Alpha-proteobacteria* became the predominant bacterial groups during the BPA biodegradation process (Yang et al., 2015). Interestingly, Peng et al.'s research found that both *Beta-proteobacteria* and *Gamma-proteobacteria* were the dominant microorganisms when TBBPA was degraded by sewage sludge, which was from an anaerobic tank of wastewater treatment facilities for TBBPA removal for several years (Wu et al., 2015). Their research concluded that both *Beta-proteobacteria* and *Gamma-proteobacteria* may biodegrade BPA, given that BPA is an important biodegradation intermediate of TBBPA (An et al., 2011).

For this study, Fig. 4 shows that there was a large variation in the proportion of *Alpha-proteobacteria* (from 56.9% to 33.3%) and *Beta-proteobacteria* (from 8.0% to 38.0%) in the microcosms during the 7-week incubation period. Other studies have reported that the genera of *Alpha-proteobacteria* can produce extracellular polymeric substances (EPS) or are motile using polar flagella, providing a protective environment for cells to grow and persist (Pang and Liu, 2007). Therefore, *Alpha-proteobacteria* appeared to be one of the major divisions in the microcosm. The ecologically diverse *Beta-*

proteobacteria were the most frequent bacteria in the clone library, consistent with published literature (Wu et al., 2015). *Beta-proteobacteria* can attach more easily to surfaces and dominated the sediment (Douterelo et al., 2013). Furthermore, this group contained many heterotrophic, chemoorganotrophic, and facultative anaerobic respiring bacteria (Haller et al., 2011). For example, members of this group, including *Dechloromonas aromatic*, were found in sediment habitats and could oxidize aromatic compounds (Coates et al., 2001). In addition, other species of this group can grow using other carbon compounds or reduced sulphur compounds (Haller et al., 2011). The sediment in microcosm was significantly polluted by different organics, including PAHs (Zhang et al., 2011). The *Beta-proteobacteria*, therefore, may be one of the major species in the microcosm. There was also many *Gamma-proteobacteria* presented in microcosm (Fig. 4); this is interesting because microorganisms from *Gamma-proteobacteria* may be linked to the removal of phenolic compounds (Yang et al., 2015), making the high presence of *Gamma-proteobacteria* expected.

Firmicutes (accounting for 12.0–34.0% of total bacteria) was the second largest phylum group in the water-sediment from the microcosm (Fig. 3). *Firmicutes* was present at 21.3% during Week 0; levels slightly decreased during Week 2, and gradually increased to 34.0% during Week 7. These changes may be that the genus *Bacillus* sp., which can degrade BPA (Li et al., 2012), is affiliated with the *Firmicutes* phylum. First, *Bacillus* sp. GZB experienced a lag phase in adapting to the water-sediment environment; levels increased as incubation continued. As such, *Firmicutes* phylum levels slightly decreased during Week 2, and then increased. *Bacteroidetes* levels increased from 8.6% to 18.6% during the incubation period, while *Actinobacteria* decreased during the incubation period.

These results demonstrated that both *Bacteroidetes* and *Bacteroidetes* may be able to degrade BPA, supporting previous studies. For example, both *Bacteroidetes* and *Actinobacteria* phyla were also present in sediment bacterial communities associated with BPA biodegradation under anaerobic conditions (Yang et al., 2015). In

addition to *Proteobacteria*, *Bacteroidetes* was the second largest group in Vidy sediment heavily contaminated with organic and inorganic pollutants (Haller et al., 2011). There was slight variation in the proportions of *Acidobacteria*, *Verrucomicrobia*, *Planctomycetes*, *Chloroflexi*, and TM7 in water-sediment from microcosm with bioaugmentation. In Yang et al.'s research, *Chloroflexi* was identified as a major bacterial group in BPA-degrading sediment (Yang et al., 2015). These results suggested that bioaugmentation or xenobiotics (BPA) could change the composition of major bacterial groups, and most of the remaining dominant bacterial groups can degrade BPA.

Bacterial community composition was further analyzed at the genus level in the clone library. There were 18 frequently detected genera in water-sediment from microcosm bioaugmented with *Bacillus* sp. GZB. As Fig. S2 shows, *Bacillus*, *Flaviumibacter*, *Dyella*, and *Thiobacillus* were the mainly genus groups in water-sediment from microcosms in this study. The relative levels of *Bacillus* genus continuously increased from 11.8% to 33.4% during the 7-week incubation period. The genus *Bacillus* was well known for its ability to degrade BPA under both aerobic and anaerobic conditions (Li et al., 2016). This made the continuous increase of the *Bacillus* genus in water-sediment from the microcosm somewhat expected. The relative levels of *Flaviumibacter* and *Thiobacillus* genus varied from 0.3% to 14.2% and from 7.5% to 5.5%, respectively. Based on these results, these genera may be able to degrade the BPA or TBBPA or other organics present in the sediment (Zhang et al., 2011; Xiong et al., 2015).

To better understand the BPA biodegradation process, a heatmap clustering analysis of the top 18 most abundant genera in each water-sediment sample was used to study microbial community compositions. As Fig. 5 shows, some genera, including *Ochrobactrum* (6.7%, *Proteobacteria*), *Bacillus* (13.4%, *Firmicutes*), *Oscillibacter* (3.4%, *Firmicutes*), *parasegetibacter* (3.8%, *Bacteroidetes*), *Cloacibacterium* (2.2%, *Bacteroidetes*), *knoellia* (7.6%, *Actinobacteria*), and TM7 (2.4%) were dominant during Week 0. As incubation time continued, *Ochrobactrum* (0.3%) became a minor genus, while




Fig. 5. Heatmap showing the top 18 most abundant genera of bacterial communities for each sample. The relative frequencies are indicated by color intensity, coded in the legend in the bottom right corner. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

Bacillus (33.4%), *Thiobacillus* (11.3%, *Proteobacteria*), *Phenyllobacterium* (14.2%, *Proteobacteria*), *Oscillibacter* (2.7%), and *Cloacibacterium* (5.5%), became the dominant genera by Week 7. There was a significant increase of *Bacillus* (from 13.4% to 33.4%) during the incubation period. There were also increases in *Thiobacillus* (from 0.7% to 11.3%) and *Phenyllobacterium* (from 0.9% to 14.2%). In summary, there was a shift in major bacterial group composition in the microcosms that were bioaugmented with *Bacillus* sp. GZB after 7 weeks of incubation.

With respect to the decrease of *Ochrobactrum* genus, previous research suggested that *Ochrobactrum* can effectively debrominate and mineralize TBBPA (An et al., 2011; Liang et al., 2016). Therefore, the decrease of *Ochrobactrum* abundance was expected, as BPA was inhibited with the increase in incubation time. Documented BPA-degrading isolates included members of *Sphingomonas*, *Pseudomonas*, *Klebsiella*, *Pandoraea*, *Alcaligenes*, *Enterobacter*, *Serratia*, *Bordetella*, *Achromobacter*, *Novosphingobium*, *Nitrosomonas*, *Cupriavidus*, *Streptomyces*, and *Bacillus* (Matsumura et al., 2009). In this study, with the exception of *Bacillus*, none of the dominant microorganisms were related to any known BPA-degraders. This suggested that *Bacillus* play an important role in BPA biodegradation in microcosms.

Previous studies have reported a link between *Thiobacillus* species and hydrogen sulfide oxidation (Gerrity et al., 2016). *Phenyllobacterium* species may be able to degrade halogenated compounds (Betancur-Corredor et al., 2015), and members of the genus *Cloacibacterium* may degrade hydrocarbon (Jurelevicius et al., 2013). However, no research has reported that these three groups can degrade BPA. Therefore, the prominent increase of *Bacillus* species in the microcosm may explain the rapid attenuation of BPA in microcosms with bioaugmentation. This suggested that bioaugmentation with *Bacillus* sp. GZB may enhance BPA biodegradation in water-sediment.

4. Conclusions

Large amounts of BPA could be biodegraded in water-sediment microcosms bioaugmented with *Bacillus* sp. GZB under aerobic conditions. *Bacillus* sp. GZB was found to play a key role in BPA biodegradation. Furthermore, a variety of relative abundant bacterial species may help increase BPA biodegradation in water-sediment microcosms with bioaugmentation. Bacterial community structure varied as incubation continued. Bioaugmentation or xenobiotics (BPA) impacted the water-sediment bacterial community structure. Finally, supplemental yeast extract, humic acid, or glucose act as electron donors/co-substrates to improve BPA biodegradation. The study might give a light to the future applications of strategy to remediate water-sediment contaminated with organics.

Acknowledgments

This work was financially supported by National Natural Science Foundation of China (41373103 and 41425015), Team Project from NSF of Guangdong Province, China (S2012030006604), and China Postdoctoral Science Foundation (2016M592548).

Appendix A. Supplementary data

Supplementary data related to this article can be found at <http://dx.doi.org/10.1016/j.chemosphere.2017.05.163>.

References

An, T.C., Zu, L., Li, G.Y., Wan, S.G., Mai, B.X., Wong, P.K., 2011. One-step process for

- debromination and aerobic mineralization of tetrabromobisphenol-A by a novel *Ochrobactrum* sp. T isolated from an e-waste recycling site. *Bioresour. Technol.* 102, 9148–9154.
- Belfroid, A., van Velzen, M., van der Horst, B., Vethaak, D., 2002. Occurrence of bisphenol A in surface water and uptake in fish: evaluation of field measurements. *Chemosphere* 49, 97–103.
- Beronius, A., Johansson, N., Ruden, C., Hanberg, A., 2013. The influence of study design and sex-differences on results from developmental neurotoxicity studies of bisphenol A, implications for toxicity testing. *Toxicology* 311, 13–26.
- Betancur-Corredor, B., Pino, N.J., Cardona, S., Peñuela, G.A., 2015. Evaluation of biostimulation and tween 80 addition for the bioremediation of long-term DDT-contaminated soil. *J. Environ. Sci.* 28, 101–109.
- Careghini, A., Mastorgio, A.F., Saponaro, S., Sezenna, E., 2015. Bisphenol A, non-ylphenols, benzophenones, and benzotriazoles in soils, groundwater, surface water, sediments, and food: a review. *Environ. Sci. Pollut. Res.* 22, 5711–5741.
- Chen, M.Y., Ike, M., Fujita, M., 2002. Acute toxicity, mutagenicity, and estrogenicity of bisphenol-A and other bisphenols. *Environ. Toxicol.* 17, 80–86.
- Coates, J.D., Chakraborty, R., Lack, J.G., O'Connor, S.M., Cole, K.A., Bender, K.S., Achenbach, L.A., 2001. Anaerobic benzene oxidation coupled to nitrate reduction in pure culture by two strains of dechloromonas. *Nature* 411, 1039–1043.
- Conte, P., Zena, A., Piliadis, G., Piccolo, A., 2001. Increased retention of polycyclic aromatic hydrocarbons in soils induced by soil treatment with humic substances. *Environ. Pollut.* 112, 27–31.
- Douterelo, I., Sharpe, R.L., Boxall, J.B., 2013. Influence of hydraulic regimes on bacterial community structure and composition in an experimental drinking water distribution system. *Water Res.* 47, 503–516.
- Eio, E.J., Kawai, M., Tsuchiya, K., Yamamoto, S., Toda, T., 2014. Biodegradation of bisphenol A by bacterial consortia. *Int. Biodegrad. Biodegr.* 96, 166–173.
- Erler, C., Novak, J., 2010. Bisphenol A exposure: human risk and health policy. *J. Pediatr. Nur.* 25, 400–407.
- Fava, F., Armenante, P.M., Kafkewitz, D., Marchetti, L., 1995. Influence of organic and inorganic growth supplements on the aerobic biodegradation of chlorobenzoic acids. *Appl. Microbiol. Biotechnol.* 43, 171–177.
- Gerrity, S., Kennelly, C., Clifford, E., Collins, G., 2016. Hydrogen sulfide oxidation in novel horizontal-flow biofilm reactors dominated by an *Acidithiobacillus* and a *Thiobacillus* species. *Environ. Technol.* 37, 2252–2264.
- Haller, L., Tonolla, M., Zopf, J., Peduzzi, R., Wildi, W., Pote, J., 2011. Composition of bacterial and archaeal communities in freshwater sediments with different contamination levels (Lake Geneva, Switzerland). *Water Res.* 45, 1213–1228.
- Hoekstra, E.J., Simoneau, C., 2013. Release of bisphenol A from polycarbonate-A review. *Crit. Rev. Food Sci. Nutr.* 53, 386–402.
- Huang, Y.Q., Wong, C.K.C., Zheng, J.S., Bouwman, H., Barra, R., Wahlstrom, B., Neretin, L., Wong, M.H., 2012. Bisphenol A (BPA) in China: a review of sources, environmental levels, and potential human health impacts. *Environ. Int.* 42, 91–99.
- Hutler Wolkowicz, I., Svartz, G.V., Aronzon, C.M., Perez Coll, C., 2016. Developmental toxicity of bisphenol A diglycidyl ether (epoxide resin badge) during the early life cycle of a native amphibian species. *Environ. Toxicol. Chem.* 35, 3031–3038.
- Jurelevicius, D., Alvarez, V.M., Marques, J.M., Lima, L., Dias, F.D., Seldin, L., 2013. Bacterial community response to petroleum hydrocarbon amendments in freshwater, marine, and hypersaline water-containing microcosms. *Appl. Environ. Microbiol.* 79, 5927–5935.
- Lee, S.W., Liu, X.S., Takeda, S., Choi, K., 2013. Genotoxic potentials and related mechanisms of bisphenol A and other bisphenol compounds: a comparison study employing chicken DT40 cells. *Chemosphere* 93, 434–440.
- Li, G.Y., Xiong, J.K., Wong, P.K., An, T.C., 2016. Enhancing tetrabromobisphenol A biodegradation in river sediment microcosms and understanding the corresponding microbial community. *Environ. Pollut.* 208, 796–802.
- Li, G.Y., Zu, L., Wong, P.K., Hui, X.P., Lu, Y., Xiong, J.K., An, T.C., 2012. Biodegradation and detoxification of bisphenol A with one newly-isolated strain *Bacillus* sp. GZB: kinetics, mechanism and estrogenic transition. *Bioresour. Technol.* 114, 224–230.
- Liang, Z.S., Li, G.Y., An, T.C., Zhang, G.X., Das, R., 2016. Draft genome sequence of a tetrabromobisphenol A-degrading strain, *Ochrobactrum* sp. T, isolated from an electronic waste recycling site. *Genome Announc.* 4 (4) <http://dx.doi.org/10.1128/genomeA.00680-16>.
- Matsumura, Y., Hosokawa, C., Sasaki-Mori, M., Akahira, A., Fukunaga, K., Ikeuchi, T., Oshiman, K.I., Tsuchido, T., 2009. Isolation and characterization of novel bisphenol-A-degrading bacteria from soils. *Biocontrol Sci.* 14, 161–169.
- Pang, C.M., Liu, W.T., 2007. Community structure analysis of reverse osmosis membrane biofilms and the significance of *Rhizobiales* bacteria in biofouling. *Environ. Sci. Technol.* 41, 4728–4734.
- Peng, Y.H., Chen, Y.J., Chang, Y.J., Shih, Y.H., 2015. Biodegradation of bisphenol A with diverse microorganisms from river sediment. *J. Hazard. Mater.* 286, 285–290.
- Quan, C., Wang, C., Duan, P., Huang, W., Yang, K., 2017. Prenatal bisphenol A exposure leads to reproductive hazards on male offspring via the Akt/mTOR and mitochondrial apoptosis pathways. *Environ. Toxicol.* 32, 1007–1023.
- Roh, H., Subramanya, N., Zhao, F., Yu, C.P., Sandt, J., Chu, K.H., 2009. Biodegradation potential of wastewater micropollutants by ammonia-oxidizing bacteria. *Chemosphere* 77, 1084–1089.
- Sun, Q., Li, Y., Chou, P.H., Peng, P.Y., Yu, C.P., 2012. Transformation of bisphenol A and alkylphenols by ammonia-oxidizing bacteria through nitration. *Environ. Sci. Technol.* 46, 4442–4448.
- Toro-Velez, A.F., Madera-Parra, C.A., Pena-Varon, M.R., Lee, W.Y., Bezares-Cruz, J.C., Walker, W.S., Cardenas-Henao, H., Quesada-Calderon, S., Garcia-Hernandez, H.,

- Lens, P.N.L., 2016. BPA and NP removal from municipal wastewater by tropical horizontal subsurface constructed wetlands. *Sci. Total Environ.* 542, 93–101.
- USEPA, 2010. Bisphenol, Action plan (CASRN 80-05-7), in: U.S. Environmental Protection Agency, 2010.
- Wang, Q., Garrity, G.M., Tiedje, J.M., Cole, J.R., 2007. Naive Bayesian classifier for rapid assignment of rRNA sequences into the new bacterial taxonomy. *Appl. Environ. Microbiol.* 73, 5261–5267.
- Wang, Z., Yang, Y.Y., Sun, W.M., Xie, S.G., 2014. Biodegradation of nonylphenol by two alphaproteobacterial strains in liquid culture and sediment microcosm. *Int. Biodeter. Biodegr.* 92, 1–5.
- Wu, Q.H., Leung, J.Y.S., Geng, X.H., Chen, S.J., Huang, X.X., Li, H.Y., Huang, Z.Y., Zhu, L.B., Chen, J.H., Lu, Y.Y., 2015. Heavy metal contamination of soil and water in the vicinity of an abandoned e-waste recycling site: implications for dissemination of heavy metals. *Sci. Total Environ.* 506–507, 217–225.
- Xiong, J.K., An, T.C., Zhang, C.S., Li, G.Y., 2015. Pollution profiles and risk assessment of PBDEs and phenolic brominated flame retardants in water environments within a typical electronic waste dismantling region. *Environ. Geochem. Health* 37, 457–473.
- Xiong, J.K., Li, G.Y., An, T.C., 2017. The microbial degradation of 2,4,6-tribromophenol (TBP) in water/sediments interface: Investigating bioaugmentation using *Bacillus* sp. GZT. *Sci. Total Environ.* 575, 573–580.
- Xiong, J.K., Li, G.Y., An, T.C., Zhang, C.S., Wei, C.H., 2016. Emission patterns and risk assessment of polybrominated diphenyl ethers and bromophenols in water and sediments from the Beiji River, South China. *Environ. Pollut.* 219, 596–603.
- Yang, L.H., Cheng, Q., Tam, N.F.Y., Lin, L., Su, W.Q., Luan, T.G., 2016. Contributions of abiotic and biotic processes to the aerobic removal of phenolic endocrine-disrupting chemicals in a simulated estuarine aquatic environment. *Environ. Sci. Technol.* 50, 4324–4334.
- Yang, Y.Y., Wang, Z., He, T., Dai, Y., Xie, S.G., 2015. Sediment bacterial communities associated with anaerobic biodegradation of bisphenol A. *Microb. Ecol.* 70, 97–104.
- Zhang, C., Zeng, G.M., Yuan, L., Yu, J., Li, J.B., Huang, G.H., Xi, B.D., Liu, H.L., 2007. Aerobic degradation of bisphenol A by *Achromobacter xylosoxidans* strain B-16 isolated from compost leachate of municipal solid waste. *Chemosphere* 68, 181–190.
- Zhang, D.L., An, T.C., Qiao, M., Loganathan, B.G., Zeng, X.Y., Sheng, G.Y., Fu, J.M., 2011. Source identification and health risk of polycyclic aromatic hydrocarbons associated with electronic dismantling in Guiyu town, South China. *J. Hazard. Mater.* 192, 1–7.
- Zu, L., Xiong, J.K., Li, G.Y., Fang, Y.J., An, T.C., 2014. Concurrent degradation of tetra-bromobisphenol A by *Ochrobactrum* sp. T under aerobic condition and estrogenic transition during these processes. *Ecotox. Environ. Safe.* 104, 220–225.